

# Acimaxyl 72 WP

## Type and composition:

Fungicide contain 64 % Mancozeb + 8% Metalaxyl w/w in the form Wettable Powder (WP) .

## Mode of Action:

Systemic preventive and curative fungicide, with contact action.

## Advantage and properties:

**Acimaxyl:** Effective to control downy mildew, early and late blight, scab, mold and other fungus disease.

**Acimaxyl:** Use on fruit trees, vegetable and other crop.

**Acimaxyl:** Mixed with common insecticides and fungicides, No mixed with alkaline material such as: Lime and sulfur, the work experience is recommended before spraying.

**Acimaxyl:** Non-phytotoxic to most plant species when used as recommended.

**Acimaxyl:** May cause allergic to skin, may affect human fertility, toxic to aquatic organism.

## Application rate:

Crop	Disease	Rate of Application/ 20L water	PHI (days)
Potatoes	late blight, early blight, tuber rot Anthracnose, leaf spot	50 g	21
Tomatoes	Late blight, early blight, spot Anthracnose, Downy mildew	70 g	3
Cucumber, Squash melon( in the ground)	Downy mildew	50 g	7
Onion, Garlic	Downy mildew, mold Blight, spotting	50 g	7
Cabbage, cauliflower Spinach	Downy mildew, spotting White rust,	50 g	7
Wheat, barely	White rust, Leaf spot	140 g/dun (50-100 L)	15
Grape, apple, peach, Pear, almond plum, banana, nectarine	Downy mildew, brown rot, scab, leaf spot Shot hole, fruit rot, Peach leaf curl	50 g	14

