



Aciprid 20% SL

Type and composition:

Insecticide contain 20 % w/v Acetamiprid in the form Soluble liquid (SL).

Mode of Action:

Systemic insecticide with translaminar activity and with contact and stomach action.

Advantage and properties:

Aciprid: effect for control many of insect such as: thrips, aphids, whitefly, leaf miner and apple fruit moth, by effect on insect central nervous system

Aciprid: Use on fruit trees and vegetable.

Aciprid: compatible with commonly used insecticides fungicides except that which have alkaline effect, preferably woks experiment initial mixing.

Aciprid: Non-phytotoxic to most plant species when used as recommended.

Aciprid: harmful if swallowed, toxic to bees, harmful to aquatic organisms.

Application rate:

| crops | pest | Rate of application/20L water | PHI |
|------------------------|-------------------------------|-------------------------------|---------|
| Cucumber, Tomato | Aphid, Thrips, Whitefly | 10 ml | 3 days |
| Eggplant, Sweet pepper | Aphid, Thrips | 5 ml | 3 days |
| Strawberry | Aphid | 5 ml | 3 days |
| Potato | Aphid | 10 ml | 7 days |
| Apple, Pear | Aphid, Fruit moth, Leaf miner | 5-10 ml | 14 days |
| Peach | Aphid, Fruit moth, Leaf miner | 5-10 ml | 7 days |
| Melon | Aphid, Trips | 5-10 ml | 3 days |
| Citrus | Aphid, Leaf miner, mealy bug | 5-10 ml | 14 days |

