



# Saltozon14% G

## Type and composition:

Insecticide contains 14 % w/w Diazinon in the form Granules (G).

## Mode of Action:

Non systemic insecticide, with contact, stomach action and respiratory system action.

## Advantage and properties:

**Saltozon:** Non systemic insecticide, with contact, stomach action and respiratory system action.

**Saltozon:** Used to control the agriculture insect such as cutworm, wireworm, and mole crickets on the vegetable

**Saltozon:** Incompatible with other pesticide.

**Saltozon:** Don't apply this product in greenhouse, and the importance distribute the pesticide equally and uniformly to get the required effectiveness of the pesticide.

**Saltozon:** Harmful if swallowed, Toxic to aquatic organism, Toxic to bees.

## Application rate:

Main crops	Typical Pests	Rate of application	Note
Cabbage, broccoli, cauliflower	Cutworm, wireworm	1.6 – 3.2 kg /D	Broadcast and incorporate into soil before planting
	Root maggots	1.6 – 2.4 kg / D	Broadcast and incorporate into soil before planting or transplanting
	Mole crickets	0.8 kg / D	Broadcast and incorporate into plant bed or field 1-2 days before sowing of seed or transplanting
Carrots	Cutworm, wireworm	1.6 – 3.2 kg/ D	Broadcast and incorporate into soil before planting
	Mole crickets	0.8 kg/ D	Broadcast and incorporate into plant bed or field 1-2 days before sowing of seed or transplanting
Lettuce	Cutworm, wireworm	1.6 – 3.2 kg/ D	Broadcast and incorporate into soil before planting
	Mole crickets	0.8 kg/ D	Broadcast and incorporate into plant bed or field 1-2 days before sowing of seed or transplanting
Watermelon muskmelon	Cutworm, wireworm	1.6 – 3.2 kg/ D	Broadcast and incorporate into soil before planting
Onion	Onion maggots, wireworm	1.6 – 3.2 kg/ D	Broadcast and incorporate into soil before planting
Radish	Mole crickets	0.8 kg/ D	Broadcast and incorporate into plant bed or field 1-2 days before sowing of seed or transplanting
Spanish	Cutworm, wireworm	1.6 – 3.2 kg/ D	Broadcast and incorporate into soil before planting
Corn	Cutworm, wireworm, seed corn maggot	1.6 – 3.2 kg/ D	Broadcast and incorporate into soil before planting
Tomato	Cutworm, wireworm	1.6 – 3.2 kg/ D	Broadcast and incorporate into soil before planting
	Mole crickets	0.8 kg/ D	Broadcast and incorporate into plant bed or field 1-2 days before sowing of seed or transplanting
Potato	Cutworm, wireworm	1.1 – 2.2 kg/ D	Broadcast and incorporate into soil before planting