

Top kill

Type and composition:

Insecticide contain 40 % w/v cypermethrin in the form Emulsifiable concentrate (EC)

Mode of Action:

Non-systemic insecticide with contact and stomach action. Also exhibits anti-feeding action. Good residual activity on treated plants.

Advantage and properties:

Top kill: Non-systemic pyrethroid insecticide, effective to control many of insect such as: cut worms, aphids, beetle, leaf miner, leaf hopper, and moth.

Top kill: Use on pome fruit, stone fruit, vegetable, field crop and ornamental plant.

Top kill: compatible with commonly used insecticides fungicides

except that which have alkaline effect, Advised to sure each case when mixing.

Top kill: Non-phytotoxic to most plant species when used as recommended.

Top kill: harmful if swallowed or inhalation, May cause allergic for respiratory system, very toxic to bees and aquatic organism.

Application rate:

Crop	pests	Rate of application	PHI
Apple, peach, apricot, almond, pear, nectarine.	Cut worms, fruit worms, beetle, aphid, sucking and chewing insect, leaf miner, leaf hopper, thrips, moths,	2-3 ml/20L water	7 days
Tomato, potato, cucumber, squash, eggplant, bean.	Aphid, thrips, caterpillar, whitefly, beetle, bugs, moths	2.5-3 ml/20L water	7 days
Cereals, sorghum, tobacco.	Aphid, locust, leaf miner, thrips, bugs, cut worms, leaf hoppers, bud worms, caterpillar,	2-3 ml/20L water	7 days
Ornamentals and forests	Aphid, leaf miners, thrips, caterpillar, beetle, bugs, moths, cut worm	2-3 ml/20L water	---

