

# Azinam

## **Type and composition:**

Fungicide contains 50 % w/v Fluazinam in the form Suspension Concentrate (SC)

## **Mode of Action:**

Fungicide with protective action has little curative or systemic activity but good residual effect.

## **Advantage and properties:**

**Azinam:** Fungicide with protective action has little curative or systemic activity but good residual effect.

**Azinam:** may be used to delay or prevent the development of resistance to single-site fungicides.

**Azinam:** will control wide range of fungal diseases like blight, rot ...etc.

**Azinam:** Compatible with other pesticide, but prefer work mini experience before mix.

**Azinam:** Non-phytotoxic to most plant species when used as recommended.

**Azinam:** harmful if inhaled, causes serious eye damage cause an allergic skin reaction, suspected of damaging fertility, very toxic to aquatic organisms, Mild toxicity to bees.

## **Application rate:**

Crop	Disease	Rate of application /20 liter water	PHI
Just seedling: Broccoli, cabbage, cauliflower, kale, mustard , turnip	Club root	Drench: 10 ml Incorporation in soil: 60 ml	50 days 20 days for mustard
Carrot	Blight, sclerotinia rot, alternaria blight	23 ml	7 days
Peas, bean, broad bean, lupine	White mold, grey mold	12-21 ml	14 day
Onion	Botrytis leaf blight, botrytis neck rot, downy mildew, purple blotch.	ml23	7 days
Potato	Late blight, white mold, powdery scab, early blight.	12-9 ml	14 days
Tomato	Botrytis, sclerotinia and late blight.	16 ml	14 days

