

Bromalone 0.1 % TP

Type and composition:

An anticoagulant Rodenticide, contain 0.1 % w/w Bromadiolone in form tracking powder.

Mode of Action:

Bromalone 0.1 % TP blocks prothrombin formation.

Advantage and properties:

Bromalone 0.1 %: Blocks prothrombin formation in liver.

Bromalone 0.1 %: Control of rats and mice (including those resistant to warfarin) in areas containing stored products, household use, industrial buildings, and other situations).

Bromalone 0.1 %: fatal if swallowed, fatal in contact with skin, fatal if inhaled, causes damage to organs through prolonged or repeated exposure, may damage the unborn child, toxic to aquatic organisms.

Application rate:

Bromalone 0.1 % TP: Control of brown rats and house mice field, black rate , roof rat in fruit orchard, vine yards, vegetable and in stored product warehouses, industrial commercial building and the inside of transport vehicles and residential areas, can prepared dry bait and distribution, then should be monitoring to control contain of active materials

Method of preparation bait

To prepare ready bait, should be mixed 1 part of bromalone 0.1 % powder with 19 part of material of bait (corn ,wheat , cereal or dry fruit with sugar) to get final concentration in bait 0.005% .

Method of use	Rate of application
At moderate infection put the bait at different dry places in application area	20-40 gm bait where the distance between it 5 m . In severe infection decreases the distance to reach 1-2 m between baits

