

Clodia 300 EC

Type and composition:

Herbicide Containing Clodinafop- Propargyl 24 % w/v+ 6% w/v Cloquintocet mexyl in form Emulsifiable concentrates (EC).

Mode of Action:

Systemic and selective herbicide, control by Fatty acid synthesis inhibitor.

Advantage and properties:

Clodia 300 EC: Systemic and selective herbicide, control by Fatty acid synthesis inhibitor.

Clodia 300 EC: Used to control of annual grasses post-emergence on wheat.

Clodia 300 EC: Incompatible with other pesticides.

Clodia 300 EC harmful if inhaled, harmful if swallowed, harmful in contact with skin, Cause an allergic skin reaction, flammable, may cause damage to organs through prolonged or repeated exposure, toxic to aquatic organism.

Application rate:

To increase the effectiveness of the pesticide must use spray oil rate of 200 ml / 20 liters of water.

Crop	Weed	Rate of application	Weed growth stage	Note
Wheat	Wild oats	13 ml / 20 L water	2 – 5 leaf	----
		17 ml / 20 L water	2 leaf to tillering	When weeds are not actively growth
		25 ml / 20 L water	-----	when mixed with broadleaves herbicide
	Phalaris spp	17 ml/ 20 L water	2 leaf to tillering	Duplicate the rate of application when mixed with broadleaves herbicide
	Annual Ryegrass	32 ml/ 20 L water	2 leaf to tillering	Duplicate the rate of application when weeds are not actively growing.

