

Index150 SC

Type and composition:

Insecticide contains 15 % w/v indoxacarb in the form suspension concentrate (SC).

Mode of Action:

Insecticide effect by contact and stomach action, work by inhibiting sodium ion entry in to nerve cells, resulting in paralysis and death of targeted pests within 2-3 days from exposure.

Advantage and properties:

Index SC: Insecticide effect by contact and stomach action, belong to oxadiazine group.

Index SC: Use on apple, pear, vegetables. Used to control Lepidoptera and other insect

Index SC: Compatible with other insecticide.

Index SC: Non-phytotoxic to most plant species when used as recommended.

Index SC: is harmful if swallowed, causes damage to organs through prolonged or repeated exposure, toxic to aquatic life, toxic to bees.

Application rate:

| Crop | Pest | Rate of application/20 L water | PHI (days) |
|--------------------------------|-------------------------------|--------------------------------|------------|
| Tomato, pepper, eggplant | leaf miner, tomato fruit worm | 5 ml | 3 days |
| Potato | Potato tuber moth | 5 ml | 3days |
| Lettuce | Worms | 5 ml | 3 days |
| Corn | Worms, corn borer | 5 ml | 3 days |
| Cabbage, broccoli, cauliflower | Diamond back moth, worms | 5 ml | 3days |
| Apple | Apple fruit worm. | 5 ml | 20 days |
| Pear | worms | 5 ml | 20 days |

