



Turan

Type and composition:

Insecticide contain 10% w/v Dinotefuran in the form soluble concentrate (SL).

Mode of Action:

Systemic insecticide with translaminar activity and with contact and stomach action.

Advantage and properties:

Turan 10% SL: Systemic insecticide with translaminar activity and with contact and stomach action.

Turan 10% SL: Use to control Aphid, thrips, leafhopper, leafminers, whiteflies.

Turan 10% SL: Use vegetables and grape.

Turan 10% SL: Compatible with other pesticide

Turan 10% SL: Non-phytotoxic to most plant species when used as recommended.

Turan 10% SL: harmful if swallowed, toxic to aquatic life, very toxic to bees.

Application rate:

Main crops	Typical pest	Rate of application/ 20 L water	PHI days
Cucumber, squash, melon, water melon	Aphid, thrips, leafhopper, leaf miners, whiteflies	10-38 ml	3
Eggplant, pepper, tomato	Aphid, leaf hoppers, leafminers, thrips, and whiteflies.	10-38 ml	3
Grapes	Mealybug , leaf hoppers, thrips.	10-28 ml	3
Broccoli, cauliflower	Aphids, leaf hoppers, whiteflies	10-38 ml	3
Celery, cress, lettuce, spinach, endive, parsley	Aphid, whitefly, leafhopper, leaf miner	10-29 ml	7
Potato	Aphid, leaf hoppers	11-14 ml	7