

Koala

Type and composition:

Fungicide contain Azoxystrobin 20% w/v + Difenconazole 12.5% w/v in the form Suspension Concentrate (SC).

Mode of Action:

Koala is protectant, curative with systemic properties fungicide, works with contact action, to eliminate many of fungal disease on wide range of agricultural crop as in the table.

Advantage and properties:

Koala: high efficiency and a double strength in the fight against fungi that consisted the persistent from other fungicides being composed of a mixture of two active ingredients in two different groups.

Koala: Use on fruit trees, vegetable.

Koala: Mixed with insecticides, and fungicides, preferably woks experiment initial mixing.

Koala: Non-phytotoxic to most plant species when used as recommended.

Koala: is harmful if inhaled, causes skin allergic, toxic to aquatic life.

Application rate:

Main crops	Target Diseases	Rate of application /20 L water	PHI
Broccoli, cabbage Cauliflower	Anthrachnose, Leaf Spot, Powdery Mildew	10-15 ml	3 days
Onion, Garlic	Botrytis Leaf Blight, Leaf Spot, Leaf Blotch, Powdery Mildew	10-15 ml	7 days
Carrots	Alternaria Leaf Blight, Leaf Spot, Powdery Mildew	10-15 ml	7 days
Chickpea	Alternaria Blight, Powdery Mildew, Rust	10-15 ml	14 days
Citrus	All type of spots, Anthracnose, Rind Blotch, Melanose, Phomopsis Stem-End Rot, Scab	18-20 ml	14 days
Cucumber, Squash Muskmelon, Watermelon	Leaf Blight, Alternaria Leaf Spot, Anthracnose, Belly Rot, Downy Mildew, Gummy Stem Blight, Canker,	10-15 ml	3 days
Grapes	Rot, Leaf Spot, Anthracnose, Downy Mildew, Leaf Blight, Powdery Mildew	10-12 ml	14 days
Peppers Eggplant	Anthrachnose, Leaf Spot, Powdery Mildew	15-10ml	3 days
Potatoes	Early Blight, Powdery Mildew, Septoria Leaf Spot	10-15 ml	14 days
Apricots, Plums Cherries, Peaches Nectarines	Alternaria Spot and Fruit Rot, Anthracnose, Brown Rot, Leaf Rust, Powdery Mildew, Scab, Shot Hole	10-15 ml	14 days
Tomatoes	Anthrachnose, Black Mold, Early Blight, Powdery Mildew, Target Spot	10-15 ml	7 days