

Comma

Type and composition:

Fungicide contains 40 % w/v Pyrimethanil in the form Suspension Concentrate (SC).

Mode of Action:

Contact fungicide with translaminar activity, providing both protective and curative efficiency, proposed inhibitor of methionine biosynthesis.

Advantage and properties:

Comma Contact fungicide with translaminar activity, providing both protective and curative efficiency, proposed inhibitor of methionine biosynthesis.

Comma: Used to control blights, rot on fruit trees, vegetable.

Comma: compatible with other pesticide except captan, but prefer work mini experience before mix.

Comma: Non-phytotoxic to most plant species when used as recommended.

Comma: very toxic to aquatic organisms.

Application rate:

Crop	Disease	Rate of application /20 liter water	PHI
Bulb vegetables	Botrytis, leaf blight, Purple blotch, Gray mold	40 ml	7 days
Grape	Gray mold	40 ml	7 days
Apricot, Nectarine, Peach, Plum	Brown rot, Blossom blight, Shot hole, Scab, Gray mold	20-40 ml	7 days
Apple	Scab	20-40 ml	14 days
Potato, Artichoke	Early blight	15 ml	7 days
Strawberry	Gray mold	20-40 ml	3 days
Tomato	Early blight, Gray mold	15 ml	3 days