

Carnival

Type and composition:

Fungicide contain Azoxystrobin 29.6 % w/v + Flutriafol 21.8 % w/v in the Form Suspension Concentrate (SC).

Mode of Action:

Azoxystrobin: fungicide with protectant, curative, eradicator, translaminar and systemic properties. Inhibits spore germination and mycelia growth, and also shows antispore activity.

Flutriafol: Contact and systemic fungicide with eradicator and protective action absorbed by foliage, with translocation acropetally in the xylem.

Advantage and properties:

Carnival: Azoxystrobin: fungicide with protectant, curative, eradicator, translaminar and systemic properties. Inhibits spore germination and mycelia growth, and also shows antispore activity.

Flutriafol: Contact and systemic fungicide with eradicator and protective action absorbed by foliage, with translocation acropetally in the xylem

Carnival: Compatible with most products.

Carnival: No phytotoxicity, when used as recommended rate.

Carnival: don't use on apple.

Carnival: harmful if swallowed, is harmful in contact with skin and is harmful if inhaled, toxic to aquatic life.

Application rate:

Crop	pests	Rate of application / 20 liter water	PHI	Note 1	Note 2
Cabbage, Cauliflower, Broccoli, mustard	Spot, Powdery Mildew, rust	7-12 ml	7	Do not apply more than 4 application per year.	Repeat as necessary if Favorable for conditions are disease development on a minimum of 7 day intervals
Tomatoes	Anthraxnose, Powdery Mildew, Black Mold, Early Blight, Late Blight, Spot		3	Do not apply more than 4 applications per year	For all other diseases apply or when preventatively for conditions are favorable disease development and throughout the continue season at a 7 – 21 day Application interval
Spinach, parsley; lettuce	Leaf Spot, Anthracnose, Downy Mildew, Powdery Rust, Mildew		7	Do not apply more than 4 applications per year.	Repeat as necessary on 7 – 14 day intervals if are favorable for conditions disease development.

Cucumber, Muskmelon, Pumpkin, Melons, Squash	Anthraxnose, Rot, Downy Mildew, Gummy Stem Blight, Leaf Spots, Powdery Mildew, Canker		3	Do not apply more than 4 applications per year	For all other diseases, apply preventatively or when conditions are favorable for disease development and throughout the continue season at 7 - 14 day intervals.
Grapes	Rot, Powdery Mildew, Downy Mildew, Spot		14	Do not apply more than 6 applications of product per acre per year.	Repeat on 14 – 21 day the intervals throughout season.
Eggplant, Pepper, Okra	Anthraxnose, Leaf Spot, Powdery Mildew		3	Do not apply more than 4 applications per year	Repeat on a 7 – 14 day interval as necessary if conditions are favorable for disease development .
Strawberry	Powdery Mildew, Anthracnose, Rot		3	Do not apply more than 4 applications per year	Repeat as necessary if favorable for conditions are disease development on a minimum of 7 day intervals.
Apricot, Nectarine, Cherry, Peach, Plum	Spot, Rot, Anthracnose, Blossom Blight, Powdery Mildew, Scab, Shot Hole, Rust		7	Do not apply more than 4 applications per year	For all other diseases apply or when preventatively for conditions are favorable for disease development and on a 7 – 14 day continue application interval.
Almond	Leaf Spot, Anthracnose, Leaf Blight, Rust, Scab, Shot Hole		14	Do not apply more than 4 applications per year	Repeat as necessary if conditions are favorable for disease development on a 7 – 21 day interval.
Corn	Rust, Leaf Blight, Spot		7	Do not apply more than 2 applications of product per growing season	For optimum results begin when disease applications as first appears and continue needed on a 7 to 10 day interval. Use
Grains	Spot, Rust, Blotch, Powdery Mildew		30	Do not apply more than 2 applications per growing season	necessary if Repeat as conditions are favorable for disease development