

# Print 40% SC

## Type and composition:

Fungicide contains 40% w/v cyazofamid in the form suspension concentrate (SC).

## Mode of Action:

Fungicide with Protective, curative Action.

## Advantage and properties:

**Print:** Foliar and soil preventive fungicide Inhibits all stage in the lifecycle of phytophthora infestans.

**Print:** Compatible with other Pesticides, prefer work mini experience before mix.

**Print:** Very Toxic to aquatic Organisms.

## Application rate:

| Crop  | Diseases   | Rate of application/20L water                             | PHI (day) | *          | **  |
|---|--|---|-----------|------------|---|
| Cabbage, broccoli cauliflower, Brussels   | Club root<br>Downy mildew  | Drench: 20-40 ml<br>Foliar: 4 ml                          | 3         | 7-10 days  | DO NOT apply more than six (1 soil + 5 foliar) applications per crop<br>DO NOT make more than three consecutive applications                      |
| Carrot  | Spot, Root Dieback (Pythium)   | 9 ml  | 14        | 21-14 days | DO NOT apply more than 5 sprays per crop  |
| Muskmelon, Watermelon<br>Pumpkin, Squash,<br>Cucumber<br>Greenhouse Cucumber                                      | Downy mildew<br>Phytophthora blight  | 3-4 ml<br>4 ml  | 3         | 7-10 days  | DO NOT apply more than six applications per crop<br>DO NOT make more than three consecutive applications<br>-Greenhouse Cucumber (4 applications) |
| Lettuce, spinach, parsley   | White rust, Downy mildew<br>Pythium Damping-off (Pythium)                                      | 4 ml  | 3         | 7-10 days  | DO NOT apply more than six applications per crop<br>DO NOT make more than three consecutive applications  |
| chickpea, lupine, bean<br>broad bean , peas   | Downy mildew<br>Phytophthora blight  | 4 ml  | 3         | 7-14 days  | DO NOT apply more than six applications per crop<br>DO NOT make more than three consecutive applications  |
| Potato, Sweet potato  | Late blight, Leaf Blight,<br>Pink Rot, Pythium Root  | Foliar:2-4 ml<br>At Planting: 9 ml<br>Lay-by/Hilling:4 ml | 7         | 7-10 days  | DO NOT apply more than 10 applications per crop<br>DO NOT make more than three consecutive applications   |
| Onion, garlic   | Downy mildew   | 4 ml  | 3         | 7-10 days  | DO NOT apply more than six applications per crop<br>DO NOT make more than three consecutive applications  |
| Eggplant, Okra, Tomato (field and greenhouse grown)<br>Tomato (Greenhouse Transplants)<br>Greenhouse Grown Pepper | Late blight, Downy mildew<br>Phytophthora blight<br>Pythium Damping-off<br>Phytophthora blight | 3-4 ml<br>4 ml<br>4ml<br>5 ml                             | 3         | 7-10 days  | DO NOT apply more than six applications per crop<br>DO NOT make more than three consecutive applications  |

\* Minimum interval between applications

\*\*Maximum number of applications