

Libra

Type and composition:

Fungicide contain Propamocarb hydrochloride 62.5% w/v and fluopicolide 6.25% w/v in the form Suspension concentrate (SC).

Mode of Action:

Systemic fungicide with prevention and curative action.

Advantage and properties:

Libra: fungicide contain **Propamocarb Hydrochloride:** systematic fungicide with productive action absorbed by the root and leaves and **Fluopicolide:** protectant fungicide with translaminar and xylem systemic movement, inhibits sporulation and mycelial growth within plant tissues.

Libra: Compatible with other insecticide and fungicide except alkaline pesticides but prefer work mini experience before mix.

Legend: harmful if swallowed, Moderate Toxic to bees, very toxic to aquatic organisms.

Application rate:

Crop	Diseases	Rate of Application /20 liter water	PHI days
Broccoli, Cabbage Cauliflower	White blister (Albugo candida) Downy mildew (Peronospora parasitica)	24-32 ml A non-ionic wetting agent should be added	3
Onion, garlic	Downy mildew (Peronospora destructor)	24-32 ml	7
Cucumber, squash Water melon, melon pumpkin	Downy mildew (Pseudoperonospora cubensis)	24-32 ml	3
Spinach ,parsley	Downy mildew (Peronospora farinosa and Peronospora parasitica)	24-32 ml	7
Lettuce	Downy mildew (Bremia lactucae)	24-32 ml	7
Potatoes	Late blight (Phytophthora infestans)	24-32 ml	14