



# Libra 68.75% SC

## Type and composition:

Fungicide contain Propamocarb hydrochloride 62.5% w/v and fluopicolide 6.25% w/v in the form Suspension concentrate (SC).

## Mode of Action:

Systemic fungicide with prevention and curative action.

## Advantage and properties:

**Libra:** fungicide contain **Propamocarb Hydrochloride:** systematic fungicide with productive action absorbed by the root and leaves and **Fluopicolide:** protectant fungicide with translaminar and xylem systemic movement, inhibits sporulation and mycelial growth within plant tissues.

**Libra:** Compatible with other insecticide and fungicide except alkaline pesticides but prefer work mini experience before mix.

**Legend:** harmful if swallowed, Moderate Toxic to bees, very toxic to aquatic organisms.

## Application rate:

Crop	Diseases	Rate of Application /20 liter water	PHI days	*	**
Broccoli, Cabbage Cauliflower	White blister (Albugo candida)  Downy mildew (Peronospora parasitica)	24-32 ml A non-ionic wetting agent should be added	3	7-10	3
Onion, garlic	Downy mildew (Peronospora destructor)	24-32 ml	7		3
Cucumber, squash Water melon, melon pumpkin	Downy mildew (Pseudoperonospora cubensis)	24-32 ml	3		2
Spinach ,parsley	Downy mildew (Peronospora farinosa and Peronospora parasitica)	24-32 ml	7		3
Lettuce	Downy mildew (Bremia lactucae)	24-32 ml	7		3
Potatoes	Late blight (Phytophthora infestans)	24-32 ml	14		2

\* Minimum interval between applications (days)

\*\*Maximum number of applications (per crop)