



Rapid

Type and composition:

Insecticide contain 48% w/v Flubendiamide in the form suspension concentrate (SC).

Mode of Action:

Insecticide with contact action.

Advantage and properties:

Rapid: Insecticide with contact action belong to diamide group for the control of various lepidopteran pests as (Diamondback moth, cabbage white butterfly).

Rapid: Compatible with other pesticide, all mixtures should be tested prior to mixing commercial quantities.

Rapid: Non-phytotoxic to most plant species when used as recommended.

Rapid: Harmful if inhaled. May irritate the eyes, Very toxic to aquatic organism, Moderate Toxic to bees.

Application rate:

Crop	Pests	Rate of application /20 liter water	PHI (Days)	Minimum Interval between Applications (Days)	Maximum Application per Growing Season
cabbage, cauliflower broccoli	Diamondback moth, cabbage butterfly Worms	2 ml	3	7-14	3
Eggplant, peppers, tomato	Worms , tomato leaf miner	2ml			
Cucumbers Melons pumpkin, squash zucchini	Worms , cucumber moth	2-3 ml			
Lettuce, cress spinach	Worms	2 ml			
potato, sweet potato, carrot radish	Diamondback moth, cabbage butterfly Worms	2 ml			
	tomato leaf miner	3 ml			
Strawberry	worms	2 ml			