

Bromac - C

Type and composition:

An anticoagulant Rodenticide, contain 0.25 % w/w Bromadiolone in form liquid.

Mode of Action:

Bromac- C 0.25% Liquid blocks prothrombin formation.

Advantage and properties:

Bromac – C 0.25 %: Blocks prothrombin formation in liver.

Bromac – C 0.25 %: Control of rats and mice (including those resistant to warfarin) in areas containing stored products, household use, industrial buildings, and other situations).

Bromac – C 0.25 % Control of brown rats and house mice field, black rate , roof rat in fruit orchard, vine yards, vegetable and in stored product warehouses, industrial commercial building and the inside of transport vehicles and residential areas,

Bromac – C 0.25 %: Use as liquid or on bait.

Bromac – C 0.25 %: Non- toxic to cats and dogs.

Bromac – C 0.25 %: is fatal if swallowed, is fatal in contact with skin, is fatal if inhaled, causes damage to organs through prolonged or repeated exposure, very toxic to aquatic life.

Application rate:

Form of application	Pest	Rate of use
Bromac – c 0.25 % liquid	Rats under hard condition like little of food and water	1 Rodenticide: 49 ready bait Or 1 Rodenticide: 49 water
Bromac –c 0.25 % as ready bait	To control rats and mice by high amount in one time at different places .	20-40 gm from bait which distributed at distance don't increase 5 m between them In severe infection the distance between bait 1-2 m





مؤسسة صناعات الكيماويات الزراعية
AGRO CHEMICALS INDUSTRIES LTD.

