

## Fyram

### Type and composition:

Fungicide contains 50 % w/v Fluopyram in the form Suspension concentrate (SC).

### Mode of Action:

Foliar fungicide with protectant action.

### Advantage and properties:

**Fyram:** Compatible with other pesticide, but prefer work mini Experiment before mix.

**Fyram:** Toxic if swallowed or absorbed through skin, harmful if inhaled, toxic to aquatic organisms.

### Application rate:

| PHI     | Application rate/20l water | Disease                                              | Crop                                          |
|---------|----------------------------|------------------------------------------------------|-----------------------------------------------|
| 3 days  | 10-5                       | powdery mildew                                       | Artichoke                                     |
|         | 10                         | Gray mold                                            |                                               |
| 3 days  | 10-5                       | powdery mildew                                       | Broccoli, kale, cabbage, cauliflower, mustard |
|         | 10-6                       | Alternaria leaf spot                                 |                                               |
|         | 10                         | Gray mold                                            |                                               |
|         | irrigation 10-5            | Alternaria, gray mold, powdery mildew, stem rot      |                                               |
| 3 days  | 10-6                       | Purple spotting                                      | Onions, garlic, potatoes, leeks               |
|         | 10                         | Leaf blight, rot                                     |                                               |
|         | 10                         | White mold                                           |                                               |
| 30 days | 10-4 irrigation            | Leaf blight, purple spot, white mold                 |                                               |
| 3 days  | 10-7                       | powdery mildew                                       | Carrot                                        |
|         | 10                         | Cottony mildew, leaf blight, leaf spot, early blight |                                               |
| 30 days | 5                          | Powdery mildew, spotting                             | Wheat, barley, oats                           |
| 7 days  | 10-7                       | Brown spot, scabies                                  | Citrus                                        |
|         | irrigation 10              | Scabies, spotting                                    |                                               |
| 14 days | 10                         | Powdery mildew, spotting                             | Corn                                          |



|        |                 |                                          |                                                     |
|--------|-----------------|------------------------------------------|-----------------------------------------------------|
| 3 days | 10-5            | powdery mildew                           | Cucumber<br>Watermelon<br>Melon<br>Pumkin<br>Squash |
|        | 10              | Leaf spot, gray mold, stem blight        |                                                     |
|        | irrigation 10-4 | Leaf spot, Fusarium wilt, powdery mildew |                                                     |
| 3 days | 10-7            | powdery mildew                           | Eggplant, peppers,<br>chili peppers<br>Okra         |
|        | 10-6            | Early blight, spotting                   |                                                     |
|        | 10              | Gray mold                                |                                                     |