





## Type and composition:

Insecticide contain Chlorantraniliprole 10% w/v + Thimamethoxam 20% w/v in the form suspension concentrate (SC).

Mode of Action:

Systemic insecticide with contact and stomach action provide residual insect control.

Advantage and properties:

**Amanda** Systemic insecticide with contact and stomach action provide residual insect control, applied to seedlings as a seedling tray or plant hole drench, highly effective in control insect because contain two active ingredients from two different groups

**Amanda**: Compatible with other pesticide, should be tested prior to mixing commercial quantities.

Amanda: Harmful if swallowed, Causes eye and skin irritation, Very toxic to bees and to aquatic organism.

Сгор	Pest	Rate of application	Planting densities / Dunum	PHI	Note
Broccoli, Cabbage Cauliflower	Aphids Cabbage Looper Cabbage Webworm Corn Earworm Diamondback Moth Thrips Whiteflies Bug	22-30 ml / 1000 seedling	4100	6 WEEKS	DO NOT apply more than 1 application of Amanda per season
Tomato, eggplant Pepper	Aphids Thrips Whiteflies Corn Earworm	37-50 ml /1000 seedling	2500		
Parsley, Spinach, Lettuce	Aphids Corn Earworm Cabbageworm Whiteflies Leafhopper and Looper Thrips	15-30ml /1000 seedling	4100	4WEEKS	

Application rate: