

TERA

Type and composition:

Insecticide contain 50 % w/w Clothianidin in the form Water Dispersible Granules (WDG).

Mode of Action:

systemic Insecticide affecting in the insect central nervous system.

Advantage and properties:

Tera: systemic Insecticide affecting in the insect central nervous system.

Tera: Used on vegetables crops, fruit trees.

Tera: Compatible with insecticides, Herbicides, Fungicide and foliar nutrient, should be tested prior to mixing commercial quantities.

Tera: harmful if inhaled, harmful in contact with skin, harmful if swallowed, may Causes eye irritation, may cause an allergic skin reaction, toxic to aquatic organisms very toxic to bees.

Application rate:

Main crops	Typical Pests	Dosage gm/20 L	PHI
Broccoli, Cabbage, Cauliflower	Aphids, Bug, Beetles, leafhoppers, leafminers, Thrips, whiteflies	2.5- 5	7 days
		35- 45 gm/donum (soil application)	21 days
Cucumber, Melons pumpkin, squash	Aphids, Bug, Beetles, leafhoppers, leafminers, Thrips, whiteflies	2.5- 5	7 days
		35-45 gm/donum (soil application)	21 days
Fig	Beetles	3-5	7 days
Gape	Bug, Beetles, leafhoppers	2.5-5	3 days
		35-45 gm/dounum (soil application)	30 days
Parsley, Spinach Lettuce, celery	Aphids, Bug, Beetles, leafhoppers, leafminers, Thrips, whiteflies	2.5-5	7 days
		35-45 gm/dounum	21 days
Peach	Aphids, Bug, leafhoppers	3-5	7 days
Apple, pear, Quince	Aphids, Bug, leafhoppers, leafminers,	3-5	7 days
Pomegranate	Aphids, Bug, leafhoppers, whiteflies	3-5	7 days
Almond	Aphids, leafhoppers, whiteflies	3-5	14 days
Potato, Sweet Potato Artichoke	Aphids, Beetles, leafhoppers, whiteflies	1.5-2.5	14 days