

Dimonil SC

Type and composition:

Fungicide contain chlorothalonil 40% w/v and Mefenoxam 4% w/v in the form Suspension Concentrate (SC).

Mode of Action:

Chlorothalonil: non-systemic foliar fungicide with protective action.

Mefenoxam: systemic fungicide with protective and curative action, taken up by leaves, stems and roots.

Advantage and properties:

Dimonil: fungicide contain **Chlorothalonil:** non-systemic foliar fungicide with protective action and **Mefenoxam:** systemic fungicide with protective and curative action, taken up by leaves, stems and roots.

Dimonil: Used to control blights, spots on vegetables.

Dimonil: Non-phytotoxic to most plant species when used as recommended.

Dimonil: Compatible with many commonly used insecticides, fungicides but prefer work mini experience before mix

Dimonil: fatal if inhaled, causes serious eye damage, may cause respiratory irritation, cause an allergic skin reaction, very toxic to aquatic life.

Application rate:

Crop	pests	Rate of application /20 liter water	PHI days
Carrot	Early blight , Late blight, spots	35-60 ml	7
Cabbage,Cauliflower Broccoli	Downy mildew, Alternaria leaf spot	35 ml	7
Cucumber, melon Muskmelon, Squash Pumpkin	Downy mildew, Anthracnose Leaf spot, Gummy stem blight, Alternaria leaf blight, scab	60-70 ml	0
Onion, Garlic Leeks	Downy mildew, Botrytis leaf blight, blotch	60 ml	7
Potato	Early blight, Late blight, rot	60 ml	14
Tomato	Early blight, Late blight, rot, leaf spot, Anthracnose	60-70 ml	5

