



# Carnival SC

## Type and composition:

Fungicide contain Azoxystrobin 29.6 % w/v + Flutriafol 21.8 % w/v in the Form Suspension Concentrate (SC).

## Mode of Action:

**Azoxystrobin:** fungicide with protectant, curative, eradicator, translaminar and systemic properties. Inhibits spore germination and mycelia growth, and also shows antispore activity.

**Flutriafol:** Contact and systemic fungicide with eradicator and protective action absorbed by foliage, with translocation acropetally in the xylem.

## Advantage and properties:

**Carnival: Azoxystrobin:** fungicide with protectant, curative, eradicator, translaminar and systemic properties. Inhibits spore germination and mycelia growth, and also shows antispore activity.

**Flutriafol:** Contact and systemic fungicide with eradicator and protective action absorbed by foliage, with translocation acropetally in the xylem

**Carnival:** Compatible with most products.

**Carnival:** No phytotoxicity, when used as recommended rate.

**Carnival:** don't use on apple.

**Carnival:** harmful if swallowed, is harmful in contact with skin and is harmful if inhaled, toxic to aquatic life.

## Application rate:

Crop	pests	Rate of application / 20 liter water	PHI	Note 1	Note 2
Cabbage, Cauliflower, Broccoli, mustard	Spot, Powdery Mildew, rust	7-12 ml	7	Do not apply more than 4 application per year.	Repeat as necessary if Favorable conditions are for disease development minimum of 7 day on a intervals
Tomatoes	Anthraxnose, Powdery Mildew, Black Mold, Early Blight, Late Blight, Spot		3	Do not apply more than 4 applications per year	For all other diseases or apply preventatively when conditions are for disease favorable development and throughout the continue season at a 7 – 21 day Application interval.



Spinach, parsley; lettuce	Leaf Spot, Anthracnose, Downy Mildew, Rust, Powdery Mildew		7	Do not apply more than 4 applications per year.	Repeat as necessary on 7 – 14 day intervals if are favorable conditions for disease development.
Cucumber, Muskmelon, Pumpkin, Melons Squash	Anthracnose, Rot, Downy Mildew, Gummy Stem Blight, Leaf Spots, Powdery Mildew Canker		3	Do not apply more than 4 applications per year	For all other diseases, apply preventatively or when conditions are for disease favorable development and throughout the continue season at 7 - 14 day intervals.
Grapes	Rot, Powdery Mildew, Downy Mildew, Spot		14	Do not apply more than 6 applications of product per acre per year.	Repeat on 14 – 21 day the intervals throughout season.
Eggplant, Pepper, Okra	Anthracnose, Leaf Spot, Powdery Mildew		3	Do not apply more than 4 applications per year	Repeat on a 7 – 14 day interval as necessary if conditions are favorable for disease development .
Strawberry	Powdery Mildew, Anthracnose, Rot		3	Do not apply more than 4 applications per year	Repeat as necessary if favorable conditions are for disease development minimum of 7 day on a intervals.
Apricot Nectarine Cherry Peach, Plum	Spot, Rot, Anthracnose, Blossom Blight, Powdery Mildew, Scab, Shot Hole, Rust		7	Do not apply more than 4 applications per year	For all other diseases or apply preventatively when conditions are for disease favorable development and on a 7 – 14 day continue application interval.
Almond	Leaf Spot, Anthracnose, Leaf Blight, Rust, Scab Shot Hole		14	Do not apply more than 4 applications per year	Repeat as necessary if conditions are favorable for disease development on a 7 – 21 day interval.
Corn	Rust, Leaf Blight, Spot		7	Do not apply more than 2 applications of product per growing season	For optimum results when begin applications disease first appears and as needed on a 7 continue to 10 day interval. Use
Grains	Spot, Rust, Blotch, Powdery Mildew		30	Do not apply more than 2 applications per growing season	necessary if Repeat as conditions are favorable disease development for