

Oxyflur 24 EC

Type and composition:

Herbicide. Containing Oxyflurofen 24 % w/v in form Emulsifiable concentrates (EC).

Mode of Action:

Selective contact herbicide, absorbed more readily by the foliage (and especially the shoots) than by the roots, with very little translocation.

Advantage and properties:

Oxyflur 24 EC: Selective contact herbicide, absorbed more readily by the foliage (and especially the shoots) than by the roots, with very little translocation.

Oxyflur 24 EC Use as per- emergence or Post-emergence control of annual broad-leaved weeds and grasses.

Oxyflur 24 EC: Preferably used with glyphosate.

Oxyflur 24 EC: harmful if inhaled and if contact the skin, Cause allergic to skin, flammable.

Application rate:

Crop	Weed	Rate of application / 20 liter water	Time of application
Cabbage, cauliflower, broccoli	control of annual broadleaved weeds and grasses	20 – 30 ml	Used in prepared land for cultivation before the transfer of seedlings
Orange, mandarin, lemon, grapefruit		40 – 50 ml	Used on newly planted trees or small, which did not bear fruit within a year, and when the length of the weed is less than 10 cm
Garlic		10 – 15 ml	When the length of garlic plant is 10 cm
Onion		10 – 15 ml	When the weed stage 2-4 leaf and plant onions in stage 3-4 real leaf
Apple, pear, almond, peach, apricot, plum, cherry, grape		30 – 40 ml	Spray directly on the soil and on the base of trees, Should non grub soil treatment because leads to reduce the effectiveness of the pesticide, Avoid contact the pesticide mist with vegetative After the swelling buds and in the configuration phase leaves.

